


A satellite image of the Earth showing the Arctic region. The image highlights the melting of sea ice and the exposure of landmasses like Greenland and parts of Europe and North America. The colors represent different land cover types: green for forests, brown for tundra or bare land, and white for ice and snow. The surrounding oceans are a deep blue.

# **The Role of Land-Cover Change in High Latitude Ecosystems: Implications for the Global Carbon Cycle**

# Global Change in High Latitude Ecosystems



# Participants

- **A. David McGuire (PI, UAF)**
- **Dave Verbyla (Co-PI, UAF)**
- **Jerry Melillo (Co-PI, MBL)**
- **Rose Meier (UAF)**
- **Dave Kicklighter (MBL)**
- **Matt Macander, Cherie Silapaswan, Aaron Woods, Qianlai Zhuang, Dorte Dissing**
- **Scott Rupp, Terry Chapin, Tony Starfield**

# **Project Goals**

- **Develop prototype spatially explicit modeling framework to elucidate how land-cover change in high latitude ecosystems influences carbon storage**
- **Apply modeling framework to assess sensitivity and uncertainty of terrestrial carbon storage responses in high latitude ecosystems**
  - **Historical responses**
  - **Future responses**

# Strategy

- **Focus on Alaska**
- **Develop historical transient land cover**
- **Develop components of modeling framework**
- **Application of modeling framework**
  - **Retrospective analyses**
  - **Prognostic analyses**

# **Why Alaska?**

- **Substantial development over last 30 years**
- **Timing and extent of disturbances known**
- **Major disturbances include fire, logging, and insect infestation**
- **Vegetation replacement may be occurring**
- **Climate change may be affecting vegetation dynamics and the frequency and extent of disturbances**
- **Process-based research (taiga/tundra LTERs)**

# **Development of Transient Land Cover**

- **Focus: 4 regions in Alaska plus Global**
  - **Tanana River Valley - fire and logging**
  - **Copper River Valley - insects and logging**
  - **Seward Peninsula - vegetation dynamics and fire**
  - **North Slope - vegetation dynamics**
  - **Global - agricultural land use**
- **Change detection**
  - **Aerial Photography**
  - **Landsat MSS, TM**
  - **AVHRR**
  - **Ancillary Data Sets**


# Study Regions in Alaska


**North Slope:  
Tundra Transition**  
Aaron Woods

**Seward Peninsula:  
Fire and Treeline Shift**  
T. Scott Rupp  
Cherie Silapaswan

**Interior: Fire  
and Logging**  
Matt Macander  
Qianlai Zhuang

**Copper River:  
Spruce Bark Beetle**  
Dave Verbyla

0 200 Kilometers  






# **Change Detection in Interior Alaska**


- **Theoretical Issues (GCP's, Grain Size, Classes)**
- **Firescar Database 1950-1997**
- **Comparisons with Landsat Imagery**
- **Detailed Analysis of Land-Cover Change:**

## **1983 Rosie Creek Fire**

- **Pre-fire Air Photos and Landsat MSS Imagery**
- **Immediate Post-fire Air Photos and Landsat TM Imagery**
- **Decadal Post-fire Air Photos and Landsat TM imagery**

# **Potential Bias in Land Cover Change Estimates Due to Positional Errors**


- **Bias can occur at any spatial scale**
- **False change may exceed real landscape change**



## Subpixel rectification error may be optimistic:


- model error varies with GCP sample size
- model error varies with transformation order

**Landsat TM 20 Classes**




10 KM

**Landsat TM 6 General Classes**





**AVHRR 20 Classes**



500 KM

**AVHRR 5 General Classes**






	20 class, 25m pixel		20 class, 1km pixel		6 class, 25m pixel		5 class, 1km pixel	
	Co-registration RMS error	Percent Change	Co-registration RMS error	Percent Change	Co-registration RMS error	Percent Change	Co-registration RMS error	Percent Change
Trial#1	1.10	7.6	0.72	10.9	1.10	8.4	0.86	1.4
Trial#2	0.86	23.7	0.89	14.3	0.73	8.2	0.87	3.5
Trial#3	1.00	8.6	0.89	20.4	0.81	10.1	0.76	2.0
Trial#4	0.75	33.2	0.95	1.8	0.94	8.5	0.96	0.1
Trial#5	0.97	5.9	1.01	13.9	0.89	15.0	0.96	0.5
Trial#6	1.06	9.7	0.94	7.1	1.02	2.2	0.97	2.7
Trial#7	0.93	8.8	0.86	6.6	1.01	10.6	0.96	1.2
Trial#8	0.71	16.1	1.01	5.4	1.09	5.8	0.99	0.6
Trial#9	0.86	4.2	1.00	10.1	1.01	4.4	0.93	1.1
Trial#10	0.90	19.8	0.82	8.4	0.87	14.2	0.96	0.9

- **Landscape metric of heterogeneity not useful; bias in landscape change varied by trial**
- **Bias in landscape change was generally more significant with heterogeneous landscapes**
- **Subpixel co-registration may not solve the problem**

# Change Detection in Interior Alaska


- Theoretical Issues (GCP's, Grain Size, Classes)
- **Firescar Database 1950-1997**
- Comparisons with Landsat Imagery
- Detailed Analysis of Land-Cover Change:
  - 1983 Rosie Creek Fire
    - Pre-fire Air Photos and Landsat MSS Imagery
    - Immediate Post-fire Air Photos and Landsat TM Imagery
    - Decadal Post-fire Air Photos and Landsat TM imagery

## Fires with Boundaries 1950-1997






**Fires with and  
without Boundaries  
1950-1997**




# Change Detection in Interior Alaska

- Theoretical Issues (GCP's, Grain Size, Classes)
- Firescar Database 1950-1997
- **Comparisons with Landsat Imagery**
- Detailed Analysis of Land-Cover Change:
  - 1983 Rosie Creek Fire
    - Pre-fire Air Photos and Landsat MSS Imagery
    - Immediate Post-fire Air Photos and Landsat TM Imagery
    - Decadal Post-fire Air Photos and Landsat TM imagery




# Interior Alaska Imagery and Firescars





# **Change Detection in Interior Alaska**

- Theoretical Issues (GCP's, Grain Size, Classes)
- Firescar Database 1950-1997
- Comparisons with Landsat Imagery
- **Detailed Analysis of Land-Cover Change:**
  - 1983 Rosie Creek Fire**
    - Pre-fire Air Photos and Landsat MSS Imagery
    - Immediate Post-fire Air Photos and Landsat TM Imagery
    - Decadal Post-fire Air Photos and Landsat TM imagery





Pre-Fire Vegetation  
1971 Air Photo Mosaic






Fire Severity  
1983 Air Photo Mosaic





Post-fire Regeneration:  
1994/95 Air Photo Mosaic









# Predicting Post-fire Regeneration in the 1983 Rosie Creek Fire




**1971 Pre-fire Vegetation**



**1983 Fire Severity**



**1994/95 Dominant Post-Fire Vegetation**



# Vegetation Regeneration after the 1983 Rosie Creek Fire (Bonanza Creek LTER)

## Developing a Predictive Model of Post-fire Regeneration Classes



- Remote sensing as primary data source: aerial photography and satellite imagery obtained before and after the fire.
- On most burned forest sites in Interior Alaska, shrub regeneration dominates a decade after the fire. What conditions are most likely to lead to alternative successional trajectories?

<b>Air Photo Class</b>	<b>Pixel Count</b>	<b>Shrub Regen. Proportion</b>
Pre-fire Hardwood	967	0.903
Pre-fire Muskeg	5304	0.609
Pre-fire Spruce	4466	0.549
Light Burn	2262	0.743
Moderate Burn	6976	0.508
Severe Burn	1499	0.888

- Topographic position, salvage logging status, and field measurements of biomass, soil, and permafrost will be used as additional explanatory variables.




Bonanza Creek Experimental Forest (1983 Rosie Creek Fire)  
Decadal Post-Fire: 1994 Air Photo and 1991 Landsat TM Imagery




# **Global Historical Agricultural Land-Use: Development of a 0.5° Boolean Data Set**

- **Based on 1-km data set developed by Ramankutty and Foley (1998)**
  - Based on version 1.2 DISCover data set
  - Based on 1992 crop cover inventory
  - Backcast to 1860 based on historical inventory
  - Aggregated to 0.5° as fractional crop cover
- **Mixture at 5° to minimize bias**
- **Time-filtered (9-years) to minimize blinking**

**Global Crop Cover in 1992 (0.5 deg resolution)  
(Ramankutty and Foley, 1998)**




## Global Crop Cover in 1992 (0.5 Degree Boolean Data Set)





# Modeling Framework for Investigating Global Change in High Latitude Ecosystems





# **Components of the Modeling Framework**


- **Physical Properties of Terrestrial Ecosystems:**  
Based on Goodrich model of permafrost dynamics
- **Physical Processes and Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Effects of Disturbance on Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Modeling the Disturbance Regime:**  
Based on physical and biological properties of ecosystems.
- **Ecosystem Structure and the Disturbance Regime:**  
Based on ALaska FRame-based EcoSystem Code (ALFRESCO)  
Starfield and Chapin. 1996. Ecol. Appl. 6:842-864
- **Modeling Vegetation Dynamics:**  
Development of Dynamic Arctic Vegetation Model (DAVM)
- **Human Dimensions:**  
Based on conceptual model of Chapin, Starfield, and Naylor

# **Components of the Modeling Framework**

- **Physical Properties of Terrestrial Ecosystems:**  
Based on Goodrich model of permafrost dynamics
- **Physical Processes and Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Effects of Disturbance on Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Modeling the Disturbance Regime:**  
Based on physical and biological properties of ecosystems.
- **Ecosystem Structure and the Disturbance Regime:**  
Based on ALaska FRame-based EcoSystem Code (ALFRESCO)  
Starfield and Chapin. 1996. Ecol. Appl. 6:842-864
- **Modeling Vegetation Dynamics:**  
Development of Dynamic Arctic Vegetation Model (DAVM)
- **Human Dimensions:**  
Based on conceptual model of Chapin, Starfield, and Naylor





# **Why is Modeling Permafrost Dynamics Important?**

- The dynamics of permafrost influences the physical environment, ecosystem function, ecosystem structure, and the disturbance regime**
- The effects of disturbance on permafrost dynamics influences vegetation trajectories after disturbance**
- Permafrost is warming in many high latitude regions**
- Areas of discontinuous permafrost are most vulnerable to melting and are also regions of higher population density**
- Thawing of permafrost has important impacts on humans in high latitude regions**




**Soil Temperature Model**

## Soil Temperature Comparison at 23 cm - Daily Output






# **Components of the Modeling Framework**

- **Physical Properties of Terrestrial Ecosystems:**  
Based on Goodrich model of permafrost dynamics
- **Physical Processes and Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Effects of Disturbance on Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Modeling the Disturbance Regime:**  
Based on physical and biological properties of ecosystems.
- **Ecosystem Structure and the Disturbance Regime:**  
Based on ALaska FRame-based EcoSystem Code (ALFRESCO)  
Starfield and Chapin. 1996. Ecol. Appl. 6:842-864
- **Modeling Vegetation Dynamics:**  
Development of Dynamic Arctic Vegetation Model (DAVM)
- **Human Dimensions:**  
Based on conceptual model of Chapin, Starfield, and Naylor






# Simulated and Observed Soil Temperature at the Northern Study Area Old Black Spruce Site (BOREAS)



Regressions represent comparisons between simulated and observed soil temperature from June 1994 to Oct. 1996.



# Simulated and Field-based Carbon Fluxes at the Northern Study Area Old Black Spruce Site (BOREAS)





Regressions represent comparisons between simulated and field-based carbon fluxes from April 1994 to Sept 1997.

# Changes in Black Spruce Carbon Storage Between 1980 and 1989 in Boreal North America

Climate based on Jones temperature anomalies,  
Hulme precipitation anomalies, and Cramer-Leeman's climatology


NEP, g C m<sup>-2</sup> yr<sup>-1</sup>




# **Components of the Modeling Framework**

- **Physical Properties of Terrestrial Ecosystems:**  
Based on Goodrich model of permafrost dynamics
- **Physical Processes and Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Effects of Disturbance on Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Modeling the Disturbance Regime:**  
Based on physical and biological properties of ecosystems.
- **Ecosystem Structure and the Disturbance Regime:**  
Based on ALaska FRame-based EcoSystem Code (ALFRESCO)  
Starfield and Chapin. 1996. Ecol. Appl. 6:842-864
- **Modeling Vegetation Dynamics:**  
Development of Dynamic Arctic Vegetation Model (DAVM)
- **Human Dimensions:**  
Based on conceptual model of Chapin, Starfield, and Naylor


## Responses of boreal forests to fire disturbance in Bonanza Creek




## Responses of boreal forests to fire disturbance in Bonanza Creek




# Soil Carbon vs. Stand age



## 1997 Soil Respiration




# Belowground Carbon vs. Stand age








## Aboveground Vegetation Carbon: TEM Simulation vs. Inventory Data for Black Spruce Forests in Alaska




## Firescars and Cohorts



# Net Biome Productivity




## Net Biome Productivity (for entire state of Alaska)



# **Simulated Carbon Flux in Alaska: Animation**

## Station BRW




For more information, visit [www.its.jhu.edu](http://www.its.jhu.edu).

# **Modeling Historical Responses of Global Terrestrial Ecosystems to changes in Atmospheric Carbon Dioxide, Climate, and Agricultural Land Use**

- **Changes in atmospheric CO<sub>2</sub> based on Keeling record.**
- **Historical temperature based on Jones et al. anomalies.**
- **Historical precipitation based on Hulme et al. anomalies.**
- **Historical land use based on boolean data sets of (1) Esser and (2) Ramankutty et al.**
- **Fate of carbon on conversion based on Houghton.**




# Global Terrestrial Carbon Flux to the Atmosphere: Effects of CO<sub>2</sub> + Climate + Landuse ( $R_H$ + conversion flux + product flux - NPP)




# **Components of the Modeling Framework**

- **Physical Properties of Terrestrial Ecosystems:**  
Based on Goodrich model of permafrost dynamics
- **Physical Processes and Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Effects of Disturbance on Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Modeling the Disturbance Regime:**  
Based on physical and biological properties of ecosystems.
- **Ecosystem Structure and the Disturbance Regime:**  
Based on ALaska FRame-based EcoSystem Code (ALFRESCO)  
Starfield and Chapin. 1996. Ecol. Appl. 6:842-864
- **Modeling Vegetation Dynamics:**  
Development of Dynamic Arctic Vegetation Model (DAVM)
- **Human Dimensions:**  
Based on conceptual model of Chapin, Starfield, and Naylor

# Annual Area Burned as a Proportion of Decadal Area Burned



# Annual Area Burned as a Proportion of Decadal Area Burned at Regional Scales in Alaska



- On average, 50-80% of the area burned in a decade is burned in the maximum fire year.
- Within smaller regions, the impact of the largest fires and fire years is greatest.
- Fire regime dominated by very large fires and fire years.

# **Evaluating the role of interannual climate variation in proportion of area burned at half-degree resolution:**

## **Logistic Regression Model:**

**Proportion Burned =  $e^{LP} / (1 + e^{LP})$  where**

$$LP = -14.939 + 0.0896T_{MAY} + 0.6478T_{JUNE} - 0.00933 \text{ SNOWPACK}_{MAY}$$

- **The model overestimates number of cells with small burns and underestimates number of cells with large burns**
- **Aggregating results to the state scale, the model explains 33.7% of the interannual variance in area burned across Alaska**
- **Over 47 years, the model predicts area burned within a few km<sup>2</sup>**


## **Evaluating the role of fire history:**

**We examined the pattern of burning in fire-history cohorts that existed in 1990:**

	<b>Burned</b>	<b>Unburned</b>
<b>Previous Burn &gt; 40 years ago</b>	<b>135</b>	<b>1675</b>
<b>Previous Burn &lt;= 40 years ago</b>	<b>54</b>	<b>1115</b>


- Of 2979 cohorts in 1990, 60.8% had not burned in last 40 years
- Of cohorts burned in 1990, 71.4% had not burned in last 40 years
- Chi-square Test:  $P = 0.0025$

# Lightning Variation







## Regional Lightning Strike Density



# Vegetation Effects





**Figure 1: Flow Chart of Tested Hypotheses**

Text explains the hypothesis tested. H-numbers refer to hypothesis.

▶ Detectable by remote sensing techniques


▽ BLM / ASF automated network data

# **Components of the Modeling Framework**

- **Physical Properties of Terrestrial Ecosystems:**  
Based on Goodrich model of permafrost dynamics
- **Physical Processes and Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Effects of Disturbance on Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Modeling the Disturbance Regime:**  
Based on physical and biological properties of ecosystems.
- **Ecosystem Structure and the Disturbance Regime:**  
Based on ALaska FRame-based EcoSystem Code (ALFRESCO)  
Starfield and Chapin. 1996. Ecol. Appl. 6:842-864
- **Modeling Vegetation Dynamics:**  
Development of Dynamic Arctic Vegetation Model (DAVM)
- **Human Dimensions:**  
Based on conceptual model of Chapin, Starfield, and Naylor

**Modeling the influence of climate variability on carbon dynamics through changes in disturbance regime and ecosystem structure and function:**

## **Coupling of ALFRESCO with TEM**




### **Simulation Specifications:**

**Spatial Scope:** 200 km by 400 km region of the Seward Peninsula

**Temporal Scope:** 1950 to 2100

**Climate Data:** Max Planck historical connected with Hadley CM2 projected climate






# **Fire History of the Seward Peninsula, Alaska Between 1950 and Present:**

## **Number of Fires and Area Burned Per Decade**


	<b>Number</b>	<b>Area (km<sup>2</sup>)</b>
<b>Observed</b>	<b>10</b>	<b>2212</b>
<b>Simulated by ALFRESCO</b>	<b>10</b>	<b>1552</b>

# ALFRESCO Simulated Fire and Vegetation Seward Peninsula AK



# **Components of the Modeling Framework**

- **Physical Properties of Terrestrial Ecosystems:**  
Based on Goodrich model of permafrost dynamics
- **Physical Processes and Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Effects of Disturbance on Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Modeling the Disturbance Regime:**  
Based on physical and biological properties of ecosystems.
- **Ecosystem Structure and the Disturbance Regime:**  
Based on ALaska FRame-based EcoSystem Code (ALFRESCO)  
Starfield and Chapin. 1996. Ecol. Appl. 6:842-864
- **Modeling Vegetation Dynamics:**  
Development of Dynamic Arctic Vegetation Model (DAVM)
- **Human Dimensions:**  
Based on conceptual model of Chapin, Starfield, and Naylor



# **Components of the Modeling Framework**


- **Physical Properties of Terrestrial Ecosystems:**  
Based on Goodrich model of permafrost dynamics
- **Physical Processes and Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Effects of Disturbance on Ecosystem Function:**  
Based on Terrestrial Ecosystem Model (TEM) - version 4.
- **Modeling the Disturbance Regime:**  
Based on physical and biological properties of ecosystems.
- **Ecosystem Structure and the Disturbance Regime:**  
Based on ALaska FRame-based EcoSystem Code (ALFRESCO)  
Starfield and Chapin. 1996. Ecol. Appl. 6:842-864
- **Modeling Vegetation Dynamics:**  
Development of Dynamic Arctic Vegetation Model (DAVM)
- **Human Dimensions:**  
Based on conceptual model of Chapin, Starfield, and Naylor



CLIMATE

ECOSYSTEM

PEOPLE



# **Significant Results**

- **Theoretical exploration of factors that influence the level of false change detection in estimates of land cover change.**
- **Making progress in developing a methodology to estimate decadal scale land-cover change in high latitude ecosystems.**
- **Progress in modeling effects of climate on permafrost dynamics, ecosystem function, and aspects of the disturbance regime.**
- **Progress in coupling models of ecosystem structure and function in high latitude ecosystems.**
- **Progress in modeling the role of elevated atmospheric CO<sub>2</sub>, climate, and agricultural land use in the global carbon cycle.**

# **Policy Implications**

- **Relevant to international negotiations with respect to controlling the concentrations of radiatively active gases in the atmosphere.**
- **Providing support for the use of spatially explicit and temporally explicit vegetation, disturbance, and climate data in the Alaska Regional Assessment.**
- **Conducting pathfinding research for the application to Alaska of the models in the Vegetation/Ecosystem Modeling and Analysis Project (VEMAP), which is an important component of the National Assessment.**
- **Regional and global analyses will contribute to the IPCC assessment process.**

# **Manuscripts in Preparation**

## **(Expect submission by end of project)**

- **Verbyla et al. Potential bias in land cover change estimates associated with positional errors**
- **Macander et al. Vegetation dynamics after fire in interior Alaska.**
- **Zhuang et al. Modeling permafrost dynamics and carbon storage in fire-disturbed ecosystems of Alaska.**
- **McGuire et al. The role of fire disturbance, climate, and atmospheric carbon dioxide in the response of historical carbon dynamics in Alaska from 1950 to 1997: A process-based analysis with the Terrestrial Ecosystem Model.**
- **McGuire et al. The role of rising atmospheric carbon dioxide, climatic variation, and agricultural land use in terrestrial carbon storage: A forward modeling comparison among four terrestrial biosphere models.**
- **Macander/McGuire et al. An analysis of the temporal and spatial dynamics of fire across Alaska between 1950 and 1997.**

# **Manuscripts in Preparation, continued...**

**(Expect submission by end of project)**

- **Dissing and Verbyla. Landscape interactions with thunderstorms in interior Alaska.**
- **Rupp et al. Historical and projected patterns of fire disturbance and vegetation dynamics on the Seward Peninsula, Alaska.**
- **McGuire et al. Historical and projected patterns of carbon storage on the Seward Peninsula, Alaska: The role of changes in atmospheric carbon dioxide, climate, simulated fire, and simulated vegetation dynamics.**
- **Rupp TS, Starfield AM, Chapin III FS (Submitted) A frame-based spatially explicit model of subarctic vegetation response to climatic change: comparison with a point model. Landscape Ecology.**
- **Rupp TS, Chapin III FS, Starfield AM (Submitted) Response of subarctic vegetation to transient climatic change on the Seward Peninsula in northwest Alaska. Global Change Biology.**

# **Next Steps**

- **Remote Sensing and Spatial Data Sets**
- **Modeling**
- **Human Dimensions**



# Next Steps

## **Remote Sensing and Spatial Data Sets:**

- Better automation of change-detection methodology
- Application of methodology to broader region  
(Alaska and beyond)
- Use of new technologies/data - MODIS, LANDSAT 7

# Next Steps

## Modeling:

- Continued development of individual modeling components
- Focused effort on a predictive model of the fire regime
- Vegetation dynamics based on both population processes and biogeochemistry
- Comparison between forward and inverse modeling approaches
- Incorporation of human dimensions

# Next Steps

## Human Dimensions:

- Fire control policy
- Direct costs of fire control
- Non-ecosystem related costs of fire
- Ecosystem-related costs and benefits of fire
- Demographic and economic conditions



**For more information look us up at**

***<http://alces.sel.uaf.edu>***